Understanding Addiction, Recovery, and Co-Occurring Mental Health Disorders

Cice Rivera Gonzalez

Touro University

Bout Me Healing

Applications of the Scholar-Practitioner Model

Journal of Psychology, Recovery & Forensic Research
Journal: ISSN: 3071-2009

Abstract

This paper focuses on understanding addiction, recovery, and co-occurring mental health disorders and how these conditions impact individuals, families, and communities. This paper, explores addiction as a complex condition influenced by biological, psychological, and environmental factors. The paper explains why this area of research was selected, based on my academic learning and professional experience in behavioral health settings. In addition, this paper discusses key challenges and opportunities in addiction and recovery research, including stigma, relapse risk, and barriers to accessing care such as financial limitations and underserved communities. This paper also examines evidence-based approaches such as mindfulness-based interventions and integrated treatment models that address mental health and substance use disorders the same time. My unique contribution to the field is presented through a holistic, integrated care approach that support emotional awareness and recovery outcomes. The paper reviews the implications of addiction research for healthcare systems, organizations, and mental health treatment, focusing on the importance of prevention, early intervention, and access to care.

Understanding Addiction, Recovery, and Co-Occurring Mental Health Disorders

Substance use disorders and mental health disorders continue to affect individuals, families, and communities throughout society, creating a long-term emotional, social, and economic challenges (Kessler et al, 2005). For example, a person struggling with substance use might experience strained family relationships, difficulty maintaining employment, and increased emotional distress, which can make recovery difficult. Substance use disorders are usually linked to conditions such as anxiety, depression, trauma, and other co-occurring mental health challenges, which can make treatment and long-term recovery more difficult (Kelly et al, 2012). For example, an individual experiencing untreated anxiety might use substances to reduce emotional distress, but this can reinforce dependency and increase relapse risk.

Research shows that addiction should not only be seen as a behavioral issue, but also as a chronic health condition that affects brain functioning, emotional regulation, and decision-making processes (McLellan et al, 2000). For example, similar to other chronic illnesses such as diabetes or hypertension, addiction might require continuous management, where individuals might experience some stability followed by relapse when support is not maintained (Volkow et al, 2016). Because of this, many researchers and healthcare professionals express the importance of integrated treatment approaches that address substance use and mental health together in treatment models (Drake et al, 2007).

For example, individuals receiving treatment for substance use disorders benefit from conditions together such as depression or anxiety in therapy, which can improve recovery and reduce relapse risk over time (Kelly et al, 2012).

One major challenge in addressing addiction and co-occurring mental health disorders is stigma, which can discourage individuals from seeking treatment and support (Hasin et al, 2013). For example, individuals with substance use disorders might delay treatment due to fear of judgment from family, employers, or healthcare providers, which can negatively impact long-term recovery. Stigma can influence how treatment systems respond to individuals, sometimes leading to delayed care instead of coordinated treatment approaches. Research shows that addressing stigma is important for improving access to treatment and encouraging intervention early in recovery (Drake et al, 2007).

Another important factor in addiction recovery is the risk of relapse, which is might be influenced by stress, emotional triggers, and untreated mental health conditions (Hendershot et al, 2011). For example, individuals in recovery might return to substance use during periods of high emotional distress or environmental pressure if coping strategies are developed. Relapse prevention strategies are a critical component of long-term recovery planning because these strategies help identify triggers, manage cravings, and maintain stability. Research shows that mindfulness-based and behavioral interventions can improve emotional regulation and reduce relapse risk by helping individuals become more aware of cravings and triggers (Bowen et al, 2014). These approaches support long-term recovery focusing on abstinence, emotional stability, and psychological stability.

Why I Selected This Area of Research

I selected addiction recovery and co-occurring mental health disorders as my area of research because of my educational background and professional experience in behavioral health settings. Throughout my career, I have worked with individuals experiencing addiction, trauma, depression, anxiety, and emotional instability in behavioral health settings. These experiences helped me observe how substance use disorders can affect different areas of a person's life, including relationships, employment, physical health, and psychological well-being (National Institute on Drug Abuse, 2020). For example, an individual struggling with addiction might experience difficulty maintaining stable employment due to attendance issues or reduced performance, which can increase stress and increase substance use patterns. Substance use disorders are usually linked to psychological distress, unresolved trauma, and chronic stress (Sinha, 2008). For example, individuals might use substances to cope with emotional pain or unresolved trauma, which can temporarily reduce distress but reinforce addictive patterns over time (Drake et al, 2007).

Research shows that addiction is influenced by stress-related neurobiological systems, which can increase vulnerability to relapse and compulsive substance use (Volkow et al, 2016). This highlights the importance of addressing psychological and biological factors in treatment. Additionally, addiction is a growing public health concern globally, with increasing rates of substance misuse, overdose, and co-occurring psychiatric condi-

tions reported throughout populations (Grant et al, 2015). These patterns show the importance of continued research and improved intervention strategies. Recovery includes abstinence, emotional stabilization, behavioral change, identity reconstruction, and long-term coping skill development (Dennis et al, 2007). For example, mindfulness-based relapse prevention has shown to improve emotional regulation and reduce relapse risk for individuals in recovery (Witkiewitz & Bowen, 2010).

Challenges and Opportunities in Addiction and Recovery Research

One major challenge in addiction and recovery research is relapse risk and difficulty maintaining long-term recovery outcomes. Another challenge in addiction is the risk of relapse, which can be influenced by stress, emotional triggers, and untreated mental health conditions (Laudet, 2008). This challenge continues to influence how individuals are treated in healthcare systems, communities, and workplaces (SAMHSA, 2023). Stigma usually discourages individuals from seeking treatment early, which can lead to severe conditions when care is accessed later. For example, individuals might avoid seeking help due to fear of judgment from family members or employers, which can delay treatment and symptoms worsen over time. Relapse is another challenge in addiction recovery. For example, individuals may return to substance use when faced with high stress or emotional instability, if coping strategies are not developed. Research shows that relapse is associated with stress exposure, environmental triggers, and untreated mental

health conditions (Moos & Moos, 2006). This shows that recovery is an ongoing process instead of a fixed outcome.

Access to care is another critical barrier, especially in underserved or rural communities where mental health and addiction treatment services are limited (Compton et al, 2007). Financial barriers, transportation limitations, and lack of specialized providers also contribute to treatment gaps. For example, individuals without health insurance or reliable transportation might struggle to attend treatment sessions or afford mental health services and addiction services. Even with these challenges, there are also important opportunities in this field. For example, one opportunity involves integrated treatment models that combine mental health care and addiction treatment have shown to improve outcomes and reduce relapse rates (Kelly et al, 2012). Additionally, mindfulness-based interventions and cognitive-behavioral approaches have shown to be effective in supporting emotional regulation and relapse prevention (Marlatt & Witkiewitz, 2005). Technology has created new opportunities for expanding access to care. For example, telehealth services, online counseling, and virtual recovery support programs help individuals to receive support in environments where traditional treatment access are limited (National Institute on Drug Abuse, 2020).

My Unique Contribution to the Field

My unique contribution to this field involves integrating psychological, emotional, and holistic perspectives into addiction recovery and mental health treatment, especially through approaches that support emotional awareness and relapse prevention strategies (Bowen et al, 2014). Mindfulness-based interventions are an area of interest in my work because this supports emotional awareness, reduce cravings, and help individuals develop healthier ways to cope with triggers and relapse risk (Bowen et al, 2014). Motivational interviewing has been shown to improve engagement in treatment and support behavioral change (Miller & Rollnick, 2013).

Co-occurring disorders require integrated care models, as untreated mental health symptoms can contribute to relapse and treatment instability (Drake et al, 2007). For example, this shows that treating substance sue and mental health together can improve and support long-term recovery outcomes (Kelly et al, 2012). My interest is in supporting approaches that address substance use and mental health at the same time instead of treating them separately. Recovery also involves identity reconstruction, emotional growth, and reintegration into society (White, 2007). Long-term recovery involves multiple ways that support sustained growth and stability (Kelly et al, 2019). For example, peer support and community-based recovery programs play an important role in sustaining long-term recovery outcomes by providing individuals with accountability, social connection, and emotional support (Dennis et al, 2007).

Implications for Organizations, Society, and Mental Health Treatment

Addiction and mental health research affects healthcare systems, organizations, and society as a whole. Integrated treatment approaches for co-occurring disorders have shown to improve long-term outcomes (Drake et al, 2007). These approaches may help reduce relapse rates by addressing substance use and mental health concerns during treatment (Kelly et al, 2012). For example, individuals receiving treatment for depression and substance use may experience better recovery outcomes than individuals receiving treatment for only one condition. Substance use disorders can affect workplace productivity, attendance, and overall employee functioning. Organizations that provide mental health support systems and employee assistance programs may improve workforce stability and employee well-being (Grant et al, 2015). For example, employees who have access to counseling or employee assistance programs may be more likely to look for support before personal challenges affect job performance.

From a societal perspective, reducing stigma can encourage individuals to seek treatment earlier (Laudet, 2008). Increasing public awareness of addiction as a chronic condition can improve treatment engagement and recovery (SAMHSA, 2023). For example, public education campaigns can help individuals recognize the signs of substance use and disorders and encourage seeking support before symptoms become severe. The criminal justice system plays an important role in addiction outcomes, as many individuals

involved in legal systems struggle with substance use and mental health disorders. Rehabilitation-focused approaches may help reduce recidivism and support long-term recovery (Volkow et al, 2016). For example, treatment and rehabilitation programs offered in correctional settings may help individuals address substance use disorders and develop healthier coping strategies before returning to the community. Public health systems continue to focus on prevention, early intervention, and expanded access to behavioral healthcare services as important strategies for addressing the addiction crisis (National Institute on Drug Abuse, 2020). For example, community outreach programs, screening, and early intervention services can help identify individuals at risk and provide access to treatment before substance use problems become severe.

In conclusion, addiction and co-occurring mental health disorders continue to create serious challenges for individuals, families, healthcare systems, and communities. Addiction is a complex condition influenced by biological, psychological, and environmental factors, and it can often occur alongside mental health conditions such as anxiety, depression, and trauma. Some challenges in addiction and recovery include stigma, relapse risk, and limited access to care. These challenges can make it harder for individuals to seek treatment and maintain long-term recovery. Addiction is a chronic condition that affects emotional regulation, stress response, and brain functioning, which can contribute to relapse risk. Effective treatment approaches include integrated care that addresses substance use and mental health at the same time. Evidence-based interventions such as mindfulness-based approaches and motivational interviewing support emotional awareness, coping skills, and behavioral change, which can improve recovery. My experience in behavioral health settings shows the importance of treating the whole person through comprehensive and integrated approaches that address mental health and substance use together. Although challenges such as stigma, relapse, and limited access to care continue to exist, research and clinical practice support the effectiveness of integrated and supportive treatment approaches. Overall, continued progress in this field can improve recovery and support healthier individuals, families, and communities.

References

American Psychiatric Association. (2022). Diagnostic and statistical manual of mental disorders (5th ed., text rev.). https://doi.org/10.1176/appi.books.9780890425787

Bowen, S., Witkiewitz, K., Clifasefi, S. L., et al. (2014). Mindfulness-based relapse prevention for substance use disorders. JAMA Psychiatry, 71(5), 547–556. https://doi.org/10.1001/jamapsychiatry.2013.4546

Compton, W. M., et al. (2007). Prevalence of substance use disorders in the United States. Archives of General Psychiatry, 64(5), 566–576. https://doi.org/10.1001/archpsyc.64.5.566

Dennis, M. L., Foss, M. A., & Scott, C. K. (2007). Recovery trajectories. Evaluation Review, 31(6), 585–612. https://doi.org/10.1177/0193841X07307771

Drake, R. E., et al. (2007). Co-occurring mental illness and substance use disorder. Psychiatric Rehabilitation Journal, 30(3), 145–155.

Grant, B. F., et al. (2015). Alcohol use disorder epidemiology. JAMA Psychiatry, 72(8), 757–766. https://doi.org/10.1001/jamapsychiatry.2015.0584

Hasin, D. S., et al. (2013). DSM-5 substance use disorders. American Journal of Psychiatry, 170(8), 834–851. https://doi.org/10.1176/appi.ajp.2013.12060782

Hendershot, C. S., et al. (2011). Relapse prevention. Substance Abuse Treatment, Prevention, and Policy, 6, 17. https://doi.org/10.1186/1747-597X-6-17

Kessler, R. C., et al. (2005). Mental disorder prevalence. Archives of General Psychiatry, 62(6), 593–602. https://doi.org/10.1001/archpsyc.62.6.593

Kelly TM, Daley DC, Douaihy AB. Treatment of substance abusing patients with comorbid psychiatric disorders. Addict Behav. 2012 Jan;37(1):11-24. doi: 10.1016/j.addbeh.2011.09.010. Epub 2011 Sep 14. PMID: 21981788; PMCID: PMC3196788.

Kelly JF, Bergman B, Hoeppner BB, Vilsaint C, White WL. Prevalence and pathways of ecovery from drug and alcohol problems in the United States population: Implications for practice, research, and policy. Drug Alcohol Depend. 2017 Dec 1;181:162-169. doi: 10.1016/j.drugalcdep.2017.09.028. Epub 2017 Oct 18. PMID: 29055821; PMCID: PMC6076174.

Laudet AB. The case for considering quality of life in addiction research and clinical practice. Addict Sci Clin Pract. 2011 Jul;6(1):44-55. PMID: 22003421; PMCID: PMC3188817.

Marlatt, G. A., & Witkiewitz, K. (2005). Relapse prevention for alcohol and drug problems: that was Zen, this is Tao. Am Psychol. 2004 May-Jun;59(4):224-35. doi: 10.1037/0003-066X.59.4.224. PMID: 15149263.

McLellan AT, Lewis DC, O'Brien CP, Kleber HD. Drug dependence, a chronic medical illness: implications for treatment, insurance, and outcomes evaluation. JAMA.2000 Oct 4;284(13):1689-95. doi: 10.1001/jama.284.13.1689. PMID: 11015800.

Miller, W. R., & Rollnick, S. (2013). Motivational Interviewing: Helping People Change (3rd ed.). Guilford Press. https://psycnet.apa.org/record/2012-17300-000

Moos RH, Moos BS. Rates and predictors of relapse after natural and treated remission from alcohol use disorders. Addiction. 2006 Feb;101(2):212-22. doi: 10.1111/j.1360-0443.2006.01310.x. PMID: 16445550; PMCID: PMC1976118.

National Institute on Drug Abuse. (2020). July 6. Treatment and Recovery. Retrieved from https://nida.nih.gov/publications/drugs-brains-behavior-science-addiction/treatment-recovery on 2026, May 29

Sinha, R. (2008). Stress and addiction. Annals of the NY Academy of Sciences, 1141, 105–130.

Substance Abuse and Mental Health Services Administration. (2023). National survey on drug use and health. https://www.samhsa.gov/data

Vollkow, N. D., et al. (2016). Brain disease model of addiction. New England Journal of Medicine, 374(4), 363–371. https://doi.org/10.1056/NEJMra1511480

White WL. Addiction recovery: its definition and conceptual boundaries. J Subst Abuse Treat. 2007 Oct;33(3):229-41. doi: 10.1016/j.jsat.2007.04.015. PMID: 17889295.

Wittkiewitz, K., & Bowen, S. (2010). Mindfulness and relapse. Journal of Consulting and Clinical Psychology, 78(3), 362–374.